

2014

# Off Center Effect on the Photoabsorption Spectrum of the Xe@C<sub>60</sub> Endohedral Fullerene

Zhifan Chen  
*Clark Atlanta University*

Alfred Z. Msezane  
*Clark Atlanta University,*

Follow this and additional works at: <http://digitalcommons.auctr.edu/caupubs>

 Part of the [Physical Sciences and Mathematics Commons](#)

---

## Recommended Citation

<http://dx.doi.org/10.1088/1742-6596/488/2/022001>.

This Article is brought to you for free and open access by the Clark Atlanta University at DigitalCommons@Robert W. Woodruff Library, Atlanta University Center. It has been accepted for inclusion in Clark Atlanta University Faculty Publications by an authorized administrator of DigitalCommons@Robert W. Woodruff Library, Atlanta University Center. For more information, please contact [cwiseman@auctr.edu](mailto:cwiseman@auctr.edu).

## Off center effect on the photoabsorption spectrum of the Xe@C<sub>60</sub> endohedral fullerene

This content has been downloaded from IOPscience. Please scroll down to see the full text.

2014 J. Phys.: Conf. Ser. 488 022001

(<http://iopscience.iop.org/1742-6596/488/2/022001>)

View [the table of contents for this issue](#), or go to the [journal homepage](#) for more

Download details:

IP Address: 144.125.40.254

This content was downloaded on 13/04/2017 at 21:20

Please note that [terms and conditions apply](#).

You may also be interested in:

[Fragmentation of positronium in collision with singly ionized lithium](#)

S Roy and C Sinha

[Fragmentation of Positronium \(Ps\) atom in collision with Li ion](#)

C Sinha and S Roy


## Off center effect on the photoabsorption spectrum of the Xe@C<sub>60</sub> endohedral fullerene

Zhifan Chen\*<sup>1</sup>, Alfred Z Msezane\*<sup>2</sup>

\* Clark Atlanta University, Atlanta GA 30314, USA

**Synopsis** DOS's of Xe atom located at different distances from the C<sub>60</sub> center have been evaluated. The results indicate that the main confinement resonances result only when the Xe atom is within about 0.3 Å around the center of C<sub>60</sub>

We have used the recently developed TDDFT method [1] to calculate the absolute differential oscillator strengths (DOS's) for the photoabsorption of the Xe atom encapsulated at different locations of the C<sub>60</sub>. This method takes three steps to evaluate the spectrum. Firstly, the structure of the C<sub>60</sub> is optimized. Secondly, the ground state eigenvalues and eigenvectors are created by solving the Kohn-Sham equation self-consistently. Thirdly the linear response of the system to the perturbation by an external electric field is described by the TDDFT. After the C<sub>60</sub> calculation a Xe atom was introduced into the C<sub>60</sub> with the locations of the Xe atom at the center and 0.2 Å, 0.3 Å, 0.4 Å, 0.5 Å, 0.8 Å, 1.5 Å, 2.0 Å away from the center. The DOS's of the Xe@C<sub>60</sub> were evaluated using the same procedure as described for the C<sub>60</sub>. Then the DOS's for the photoabsorption of the encaged Xe atom were extracted by subtracting the DOS's of the fullerene from the corresponding DOS's of the Xe@C<sub>60</sub>.




**Figure 1.** Photoabsorption spectrum from bottom up, Xe at the center, 0.2 Å, 0.3 Å and 0.4 Å away from center, respectively

<sup>1</sup>E-mail: [zchen@cau.edu](mailto:zchen@cau.edu)

<sup>2</sup>E-mail: [amsezane@cau.edu](mailto:amsezane@cau.edu)

Fig. 1 demonstrate that the main confinement resonances result only when the Xe atom is located in a very small region of the C<sub>60</sub>. This region is about 0.3 Å around the center of the C<sub>60</sub> (3.5 Å is the C<sub>60</sub> radius). These results also explain the absence of the confinement resonances in the photoionization of the encapsulated lanthanide atoms such as Ce@C<sub>82</sub> [2], as these atoms (Ce, Pr) are usually located 1.8-2.0 Å off the center of the C<sub>82</sub>.



**Figure 2.** Solid (red) curve is the DOS's of free Xe atom, dashed (blue) curve represents the DOS's of Xe located 2 Å away from the center of C<sub>60</sub>

Fig. 2 indicates that the DOS's of the Xe atom located far away from the center is almost the same as that of the free Xe atom

This work was supported by U. S. DOE, AFOSR and Army Research.

### References

- [1] Zhifan Chen and A Z Msezane 2012 *Phys. Rev. A* **86** 063405
- [2] A Müller *et al* 2008 *Phys. Rev. Lett.* **101** 133001

