

2015

Charge Modification of Metal Atoms: Catalysis of Water to Peroxide

Kelvin Suggs
Clark Atlanta University

Filmon Kiros
Clark Atlanta University

Aaron Tesfamichael
Clark Atlanta University

Zineb Felfli
Clark Atlanta University

Alfred Z. Msezane
Clark Atlanta University

Follow this and additional works at: <http://digitalcommons.auctr.edu/caupubs>

 Part of the [Cellular and Molecular Physiology Commons](#)

Recommended Citation

Suggs, Kelvin; Kiros, Filmon; Tesfamichael, Aaron; Felfli, Zineb; and Msezane, Alfred Z., "Charge Modification of Metal Atoms: Catalysis of Water to Peroxide" (2015). *Clark Atlanta University Faculty Publications*. 39.
<http://digitalcommons.auctr.edu/caupubs/39>

This Article is brought to you for free and open access by the Clark Atlanta University at DigitalCommons@Robert W. Woodruff Library, Atlanta University Center. It has been accepted for inclusion in Clark Atlanta University Faculty Publications by an authorized administrator of DigitalCommons@Robert W. Woodruff Library, Atlanta University Center. For more information, please contact cwiseman@auctr.edu.

Charge modification of metal atoms: Catalysis of Water to Peroxide

This content has been downloaded from IOPscience. Please scroll down to see the full text.

2015 J. Phys.: Conf. Ser. 635 052018

(<http://iopscience.iop.org/1742-6596/635/5/052018>)

View [the table of contents for this issue](#), or go to the [journal homepage](#) for more

Download details:

IP Address: 144.125.40.254

This content was downloaded on 20/04/2017 at 19:43

Please note that [terms and conditions apply](#).

You may also be interested in:

[Material Research on High-Quality Passivation Layers with Controlled Fixed Charge for Crystalline Silicon Solar Cells](#)

Tomihisa Tachibana, Takashi Sameshima, Yuta Iwashita et al.

[Polarity in oxide ultrathin films](#)

Claudine Noguera and Jacek Goniakowski

[Scanning gate microscopy of magnetic focusing in graphene devices: quantum versus classical simulation](#)

M D Petrovi, S P Milovanovi and F M Peeters

[X-ray fluorescence yields for light emitter atoms: fluorine](#)

E Hartmann and R Der

[Insulator–metal transition of VO₂ ultrathin films on silicon: evidence for an electronic origin by infrared spectroscopy](#)

W W Peng, G Niu, R Tétot et al.

[Origin of valence and core excitations in LiFePO₄ and FePO₄](#)

M

K Kinyanjui, P Axmann, M Wohlfahrt-Mehrens et al.

[Recoil-induced electronic excitation and ionization in one- and two-electron ions](#)

L Wauters, N Vaeck, M Godefroid et al.

[First principles studies of complex oxide surfaces and interfaces](#)

Claudine Noguera, Fabio Finocchi and Jacek Goniakowski

Charge modification of metal atoms: Catalysis of Water to Peroxide

Kelvin Suggs^{*‡}, Filmon Kiros^{*}, Aaron Tesfamichael[‡], Zineb Felfli^{*2} and Alfred Z. Msezane^{*1}

^{*}Center for Theoretical Studies of Physical Systems, Clark Atlanta University, Atlanta, Georgia 30314, USA

[‡]Department of Chemistry, Clark Atlanta University, Atlanta, Georgia 30314, USA

Synopsis Here we investigate the optimization of the catalytic behavior of Au, Ag, Pd, Rh, and Os atomic systems via charge modification and conclude that anionic atomic systems are optimal and therefore ideal for catalyzing the oxidation of water to peroxide, with anionic Os being the best candidate. This conclusion should also be applicable to the oxidation of methane to methanol without CO₂ emission, with the anionic Os being the best catalyst.

In [1] it was demonstrated that the fundamental mechanism of negative ion catalysis of water to peroxide using atomic Au⁻ is the formation of the anionic Au⁻(H₂O)_{1,2} molecular complex during the transition state, which breaks up the H-O bonds, thereby promoting the formation of the H₂O₂ in the presence of O₂. The relatively large electron affinity (EA) of Au contributes to its anionic catalytic effectiveness. Our complex angular momentum (CAM) method [2] has been used to select the above metal atoms through the accurate determination of their large EAs for the investigations. Density Functional Theory (DFT) and dispersion corrected DFT approaches have been employed for the transition state

calculations. Geometry optimization of the structural molecular conformation utilized the gradient-corrected Perdew-Burke-Ernzerhof parameterizations [3] of the exchange-correlation as implemented in DMol3 [4]. A tolerance of 1x10⁻³ Ha was used with a smearing value of 0.005Ha.

Transition state calculated results, given in Fig. 1 show the catalyzing effects of cationic, neutral, and anionic Ag, Au, Pd, Rh, and Os atoms when synthesizing water to peroxide. Clearly, the Au⁺ is the least efficient of the given systems as shown by a barrier of ~1.3eV. Conversely, Os⁻ is energetically the most favored with a barrier of ~0.16eV as shown in Fig. 1(b).




Figure 1 (a) and (b): Potential energy transition state barrier calculations of cationic, neutral, and anionic Au, Ag, and Pd and Rh, and Os represented by orange, red, and blue and olive and black, respectively. The standard comparative reaction energy without catalysis is represented by the purple line.

The discovery that cationic systems increase the transition energy barrier in the synthesis of peroxide could render them important as inhibitors in controlling and regulating catalysis. These findings usher in a fundamental and comprehensive atomic theoretical framework for the generation of tunable catalytic systems. We conclude that anionic atomic systems are optimal for catalyzing the oxidation of water to peroxide, with Os⁻ being the preferred candidate among the atomic ions proposed in this work. Whether this is consistent across interdisciplinary systems, reaction types, and atomic cluster substrates is yet to be determined.

¹E-mail: amsezane@cau.edu

²E-mail: zfeldli@cau.edu

The structure and dynamics of these atoms with relatively large EAs will facilitate the understanding of the recently synthesized Ag@Au@Pt nanosystem [5] with greatly enhanced plasmonic properties as well as lead to the creation of tunable supercatalysts.

References

- [1] A. Tesfamichael *et al.* 2013 JNR 15 1333.
- [2] D. Sokolovski *et al.* 2007 Phys. Rev. A **76** 012705
- [3] A. Tkatchenko, and M. Scheffler 2009 Phys. Rev. Lett. **102** 73005.
- [4] DMol3 2011 Accelrys Software Inc.: San Diego, CA
- [5] N. Deogratias *et al.* 2015 Nano Res. **8** (1) 271

