

January 1941

Some reactions of p-methoxy-diphenylbutadiene

Joe Marion Hill
Atlanta University

Follow this and additional works at: <http://digitalcommons.auctr.edu/dissertations>

Recommended Citation

Hill, Joe Marion, "Some reactions of p-methoxy-diphenylbutadiene" (1941). *ETD Collection for AUC Robert W. Woodruff Library*. Paper EP17400.

This Dissertation/Thesis is brought to you for free and open access by DigitalCommons@Robert W. Woodruff Library, Atlanta University Center. It has been accepted for inclusion in ETD Collection for AUC Robert W. Woodruff Library by an authorized administrator of DigitalCommons@Robert W. Woodruff Library, Atlanta University Center. For more information, please contact cwiseman@auctr.edu.

INFORMATION TO USERS

This manuscript has been reproduced from the microfilm master. UMI films the text directly from the original or copy submitted. Thus, some thesis and dissertation copies are in typewriter face, while others may be from any type of computer printer.

The quality of this reproduction is dependent upon the quality of the copy submitted. Broken or indistinct print, colored or poor quality illustrations and photographs, print bleedthrough, substandard margins, and improper alignment can adversely affect reproduction.

In the unlikely event that the author did not send UMI a complete manuscript and there are missing pages, these will be noted. Also, if unauthorized copyright material had to be removed, a note will indicate the deletion.

Oversize materials (e.g., maps, drawings, charts) are reproduced by sectioning the original, beginning at the upper left-hand corner and continuing from left to right in equal sections with small overlaps.

ProQuest Information and Learning
300 North Zeeb Road, Ann Arbor, MI 48106-1346 USA
800-521-0600

UMI[®]

NOTE TO USERS

This reproduction is the best copy available.

UMI[®]

SOME REACTIONS OF
P-METHOXY-DIPHENYLBUTADIENE

A THESIS
SUBMITTED TO THE FACULTY OF ATLANTA UNIVERSITY
IN PARTIAL FULFILLMENT OF THE REQUIREMENTS FOR
THE DEGREE OF MASTER OF SCIENCE

BY
JOE MARION HILL

DEPARTMENT OF CHEMISTRY

ATLANTA, GEORGIA

JUNE 1941

UMI Number: EP17400

UMI[®]

UMI Microform EP17400

Copyright 2007 by ProQuest Information and Learning Company.
All rights reserved. This microform edition is protected against
unauthorized copying under Title 17, United States Code.

ProQuest Information and Learning Company
300 North Zeeb Road
P.O. Box 1346
Ann Arbor, MI 48106-1346

TABLE OF CONTENTS

ii


Chapter	Page
I THEORETICAL	1
II EXPERIMENTAL	9
III SUMMARY	14
IV BIBLIOGRAPHY	15

CHAPTER I


THEORETICAL

Ever since Thiele introduced his theory of addition to conjugated unsaturated compounds, numerous investigators have pointed out that it does not cover all the experimental facts.¹


In brief, Thiele postulated residual valencies² resulting from the excess of affinity left from the union of carbon to carbon:



Due to the close proximity of the 2,3-carbon atoms the residual valencies neutralize each other leaving only the 1,4-positions free.



Thus any addenda to the conjugated compound would give the 1,4-addition product. As the addenda would require more affinity than that offered by the residual valencies, the double bonds would be ruptured and a new double bond would form at the 2,3-position.³



It has been shown that universal 1,4-addition to conjugation,

¹Henry Gilman, "Organic Chemistry," John Wiley and Sons, Inc., New York, 1938, I, 575-576.

²J. B. Cohen, "Organic Chemistry for Advanced Students," Longmans, Green, and Co., New York, I, 145.

³Henry Gilman, op. cit.




as Thiele postulated, does not occur;¹ consequently much experimentation has been done to show the true nature of addition to conjugated compounds.

This investigation was undertaken in order to determine the position of addition of bromine to 1-para-methoxy-phenyl-4-phenylbutadiene.

Ingold has predicted that compounds of this nature would add bromine in the 3,4-position;² so that a brief resume of his theory will not be out of place.

Ingold⁵ divided addition to unsaturated compounds into three classes: (1) both parts of the compound added are stable anions: e.g., Cl₂, Br₂, OHCl, etc., which may enhance or retard anionotropic mobility; (2) an anion-cation grouping: e.g., HCl, HBr, HCN, etc., which may also enhance or retard anionotropic mobility; and (3) a cation-cation grouping: e.g., H₂.

Starting from the demonstration that one atom of halogen adds initially,⁴ Ingold⁵ postulates the following series of reactions for butadiene:



¹Straus, *Ber.*, 42, 2866 (1909).

²Ingold and Smith, *J. Chem. Soc.*, 2753 (1931).


³Ingold and Burton, *J. Chem. Soc.*, 904 (1928).

⁴Francis, *J. Am. Chem. Soc.*, 47, 2344 (1925); Terry and Eichelberger, *J. Am. Chem. Soc.*, 47, 1067 (1925).

⁵Ingold and Burton, *op. cit.*


The part of the molecule of (II) under the bracket is the anionotropic system which is activated by the CH_2Br -grouping to give the 1,2 and 1,4 ions. As CH_2Br - has just about the same effect as hydrogen as an activating agent, both the 1,2 and the 1,4-dibromides of butadiene should be formed. Farmer and co-workers¹ have isolated and characterized the two dibromides.

Phenylbutadiene² adds chlorine in the 3,4-position in accordance with Ingold's prediction.³



The phenyl group, a stronger electronegative group than the CH_2Cl -group, prevents the anionotropic shift; therefore the 3,4 dichloride is formed almost exclusively.

Ingold⁴ has also predicted that conjugated substituted diphenylbutadiene compounds with substituents having unshared pairs of electrons should add bromine in the 3,4-position, because the unshared electrons may be displaced consecutively throughout the conjugated system to build up a region of high electron density on the fourth carbon atom. For example, 1-para-bromophenyl-4-phenylbutadiene would react in this manner:



The bromophenyl-group, being a more electronegative group than the phenyl group, prevents the anionotropic shift; and only

¹Farmer, Lawrence, and Thorpe, *J. Chem. Soc.*, 729 (1928).

²Muskat and Huggins, *J. Am. Chem. Soc.*, 51, 2496 (1929).


³Ingold and Burton, *op. cit.*

⁴Ingold and Smith, *op. cit.*

the 3,4-dibromide will be formed. Yokely¹ has shown that para-bromodiphenylbutadiene does add bromine in the 3,4-position.


Proctor,² working with the corresponding para-chloro compound, got a 3,4-dichloride. This result is also in complete accord with Ingold's theory.

Similarly, 1-para-methoxy diphenylbutadiene should add bromine in the 3,4-position:




Para-methoxy diphenylbutadiene was prepared by the method of Meerwein³ and also by the method of Kuhn and Winterstein.⁴

Meerwein's method is based upon the fact that aromatic diazo compounds will react with α,β -unsaturated carbonyl compounds to give unsaturated hydrocarbons.



The lactone, formed as an intermediate,⁵ breaks down, loses carbon dioxide, and forms the butadiene.



This method gave 16% yields of the para-methoxy compound as compared to 25% yields of the para-nitro-diphenylbutadiene and 40% yields of the substituted stilbenes prepared by Bergmann

¹Yokely, "Studies in the Butadiene Series: Addition of Bromine and Hydrogen Bromide to 1-para-Bromo-phenyl-4-phenylbutadiene," Unpublished Master's Thesis, Department of Chemistry, Atlanta University, 1939.

²Proctor, "The Preparation and Chlorination of 1-para-Chloro-phenyl-4-phenylbutadiene," Unpublished Master's Thesis, Department of Chemistry, Atlanta University, 1932.

³Meerwein, Buchner, and Emster, *J. Prakt. Chem.*, **152**, 237 (1939).

⁴Robinson, "Organic Synthesis," John Wiley and Sons, New York, 1935.

⁵Meerwein, Buchner, and Emster, *op. cit.*

and Weinberg¹ and Meerwein and co-workers² respectively. The product is difficult to purify, as a great deal of tarry material along with some dye is formed as a by-product of this reaction.

Kuhn and Winterstein's³ method gave a product which was easily purified, but, on account of difficulties experienced in obtaining enough para-methoxyphenylacetic acid to prepare workable quantities of the butadiene, the method was abandoned. By condensing a small quantity of para-methoxyphenylacetic acid with cinnamic aldehyde,⁴ a product was obtained which was identical with the product obtained by Meerwein's method. When the two products were mixed they gave the same melting point.

To secure evidence of the structure of the butadiene a sample was condensed with maleic anhydride in benzene. Some difficulty was experienced in obtaining a product as condensation took place only after refluxing for sixteen hours.

The maleic anhydride condensation, the so called diene synthesis of Diels and Alder,⁵ has been accepted as a means of identification of conjugated compounds, and, as such, it is used here. Occasionally, however, maleic anhydride will condense with compounds other than the butadienes.

Besides maleic anhydride, maleic acid, acrolein, or naphthaquinone might have been used, the condensation usually taking place in the 1,4-position.


¹Borgman and Weinberg, J. Org. Chem., 6, 134 (1941).

²Meerwein, Buchner, and Emster, op. cit.

³Robinson, op. cit.

⁴Ibid.


⁵Allen, J. Chem. Ed., 10, 494 (1933).



When a sample of the diene was saturated with bromine, a fine white powder was isolated from the reaction mixture. Upon repeated recrystallization, this powder melted at 190° with much decomposition. The analyses for bromine were higher than the calculated value for the tetrabromide of para-methoxy-diphenylbutadiene.

To gain further knowledge of the structure of the butadiene prepared, a portion of it was treated with an excess of ozone; and the solution resulting from the hydrolysis of the ozonide was analyzed qualitatively for the fragments.

The decomposed ozonide should give benzaldehyde, para-methoxy benzaldehyde, and glyoxal or the corresponding acids.



Oxalic acid was identified; and, after oxidation of an oily residue, para-methoxybenzoic acid was isolated. No benzoic acid or benzaldehyde was found.

The result of the ozonolysis, along with the maleic anhydride derivative, gives evidence of the conjugated structure of para-methoxy-diphenylbutadiene.

The oil resulting from this reaction was treated with hydrogen peroxide with no visible effect. The fragments expected from the oxidation of the dibromide were not obtained. This is probably due to the instability of the dibromide.

The literature records the fact that the methoxycinnamic acids and esters react abnormally with bromine, giving substitution products instead of addition products. Davies and Davies,¹ working with the meta-methoxycinnamic acid derivatives; Hanson and Williams,² using the ortho-methoxy- and the para-methoxycinnamic acids; and Riemer and Tobin,³ working with the 2,4-dimethoxycinnamic acids, all obtained large amounts of substitution products, along with the addition products, when these compounds were treated with halogen.

Some time ago, Straus,⁴ working with similar acids, found that the highly activating methoxy group fostered substitution instead of addition. He noted that the amount of the substitution products was increased by the position and number of methoxy groups. In many cases he isolated the substitution product.

In view of Straus' work, it would be expected that paramethoxy diphenylbutadiene would react with bromine by substitution more readily than by addition. The experimental results, while far from complete, apparently support this conclusion.

The prediction that bromine would add 1,2 to paramethoxydiphenylbutadiene⁵ could not be verified as the dibromide was unstable.

¹Davies and Davies, *J. Chem. Soc.*, 602 (1928).

²Hanson and Williams, *J. Chem. Soc.*, 1059 (1930).

³Riemer and Tobin, *J. Am. Chem. Soc.*, 52, 341 (1930).

⁴*Ibid.*

⁵Ingold and Smith, *op. cit.*

CHAPTER II

EXPERIMENTAL

Preparation of 1-para-Methoxyphenyl-4-Phenylbutadiene. --

The butadiene was prepared after the manner of Meerwein.¹ The cinnarylidene acetic acid used was prepared according to Riiber's method.²

Para-anisidine (44 g.) was suspended in 200 cc. of 30% hydrochloric acid and diazotized at 0° with sodium nitrite (18 g.) in 40 cc. of water. The diazo solution was added dropwise to the cinnarylidene acetic acid (32 g.) suspended in 350 cc. of acetone which had been cooled to 5°. Then cupric chloride (10.6 g.) and sodium acetate (55 g.) in 50 cc. of water were added. Nitrogen was evolved at once, and the solution developed a deep purple color. After stirring for one hour at 15°, the reaction mixture was steam distilled until the distillate became clear. The solution was filtered while hot, and the non-volatile tarry residue was dissolved in hot glacial acetic acid. The solution was filtered while hot; and, when cool, the precipitated diene was filtered off. The precipitate was recrystallized several times from an alcohol-benzene mixture. The product isolated melted at 159°.

Anal. Calcd. for $C_{17}H_{16}O$: C, 86.4, H, 6.8. Found: C, 86.6, 86.7, 86.6; H, 7.1, 7.3, 7.0.

The butadiene was also prepared by the method of Kuhn and Winterstein.³ The para-methoxy-phenylacetic acid used was

¹Meerwein, Buchner, and Emster, J. Prakt. Chem., 152, 237 (1939).

²Riiber, Ber., 37, 2272 (1904).

³Robinson, op. cit.

prepared by the method of Cain, Simonsen, and Smith.¹

Para-methoxy-phenylacetic acid (11 g.), cinnamic aldehyde (9 g.), acetic anhydride (10 cc.), and lead monoxide (7.5 g.) were refluxed together for five hours in a flask protected from atmospheric moisture by a calcium chloride tube. The dark, hot solution was poured into a beaker and allowed to stand overnight. The grayish mass was moistened with alcohol, filtered, and recrystallized twice from toluene. A mixture of this product and the substance prepared by Moorwain's method melted at 159°.

Condensation with Maleic Anhydride. -- Maleic anhydride (0.41 g.) was dissolved in 35 cc. of benzene and filtered into a long neck flask containing para-methoxy-diphenylbutadiene. The mixture was refluxed for sixteen hours, concentrated, and the precipitate was recrystallized three times from toluene. The pure product melted at 139°. The yield was almost quantitative.

Anal. Calcd. for $C_{21}H_{18}O_4$: C, 75.4; H, 5.4. Found: C, 75.0, 73.3; H, 5.3, 3.7.

Attempted Preparation of the Tetrabromide of para-Methoxy-Diphenylbutadiene. -- A sample (1.5 g.) of the diene was dissolved in chloroform, cooled to 0°, and bromine was added in excess. Ligroin (b.p. 65°-70°) was added, and, after standing for one hour, the solution was filtered. After recrystallization several times from ligroin, a white powder melting at 190° with much decomposition was obtained. The yield of the recrystallized product was small.

The results of the analyses are approximately the same as the calculated value for the percent of bromine in para-bromo-diphenylbutadiene.

¹Cain, Simonsen, and Smith, J. Chem. Soc., 103, 1035 (1935).

Anal. Calcd for $C_{17}H_{16}OBr_4$: Br, 57.5. Found: Br, 64.8, 65.0, 65.4.

Anal. Calcd for $C_{16}H_{13}Br_5$: Br, 66.4

Ozonolysis of para-Methoxy-Diphenylbutadiene. -- A sample of the butadiene was dissolved in chloroform and ozone was passed through the solution which was cooled in a salt-ice bath. After eight hours of ozonization, the chloroform was removed by suction, and 5 cc. of dilute potassium hydroxide in 25 cc. of water were added; the mixture was warmed on a water bath for one hour.

The alkaline solution was extracted several times with ether, the extract was dried over sodium sulfate (anhydrous), and the ether was removed by suction. The oil which remained was dissolved in a slight amount of alcohol and 2 cc. of dilute potassium hydroxide and 2 cc. of 30 % hydrogen peroxide were added. The solution was warmed on the water bath for fifteen minutes, and the alcohol was removed by evaporation.

The alkaline mixture was acidified, and the white precipitate was recrystallized from water and from dilute alcohol. The fluffy solid melted at 184° . Mixing the unknown with a known sample of para-methoxy-benzoic acid caused no change in the melting point of the known acid.

The alkaline solution from the decomposition of the ozonide was acidified with sulfuric acid, made alkaline with ammonium hydroxide, and then acid with acetic acid. Boric acid (2 g.) was added. After warming and stirring to dissolve the solid, 5 cc. of calcium chloride solution were added, and the solution was set aside for ten minutes with occasional stirring. Then the solution was heated to boiling, filtered, and the precipitate was washed with hot water until all the reducing material was removed.

Five cc. of hot dilute sulfuric acid were poured on the filter paper, and the filtrate was collected in a test tube. A drop of dilute potassium permanganate solution was bleached instantly by the hot acid solution. This proved the presence of oxalic acid.

No other substances were isolated from the ozonization.

Attempted Preparation of the Dibromide of para-Methoxy-Diphenylbutadiene. -- In the dark, bromine (3.36 g.) dissolved in 25 cc. of chloroform was added dropwise through a separatory funnel to a chloroform solution of the butadiene (5 g.) protected by a calcium chloride tube. The butadiene solution was kept at 0° through-out the addition of bromine after which it was allowed to stand over night. The solution darkened, and upon removal of the solvent a black residue was obtained. Repeated recrystallizations from chloroform by the addition of ligroin (b.p. 65°-70°) yielded a small amount of tan powder melting at 123°-130° with decomposition. The analyses showed that the powder was not the dibromide.

Anal. Calcd. for $C_{17}H_{16}OBr_2$: Br, 40.4. Found: Br, 47.5, 45.9, 45.6.

Another sample was treated with bromine in the same manner as before with the exception that the chloroform was removed immediately with the aid of an oil pump. Repeated recrystallization of the black residue yielded a small amount of the original butadiene.

To another sample, slightly more than the calculated amount of bromine was added, and the cold solution was ozonized directly for eight hours. After removing the solvent by suction, the oily ozonide was decomposed by adding 25 cc. of water and by

warming on a water bath. The water solution was extracted several times with ether, and the extract was dried over anhydrous sodium sulfate. The oil left after the removal of the ether was dissolved in dilute alcohol, and 2 cc. of 30% hydrogen peroxide were added. The oil showed no visible change after the oxidation and was not analyzed.

Another portion of the diene (1.2876 g.) was brominated in the manner stated, and an equal volume of cold acetone was added to the chloroform solution. After the method of Farmer and Scott,¹ potassium permanganate (3.932 g.) and magnesium sulfate (3.932 g.), dissolved in 60 cc. of water, were added dropwise with vigorous shaking to the chloroform-acetone solution. The mud formed was filtered, and the acetone was removed by bubbling a rapid stream of air through the solution. The chloroform was removed by evaporation. The oily material obtained here was combined with the material isolated from the mud.

The mud was suspended in water acidified with sulfuric acid, and enough sodium bisulfite to bring the mud into solution was added. The acid mixture was extracted with ether and the extract neutralized with a few drops of saturated sodium bicarbonate solution. Then the ether was dried over anhydrous sodium sulfate and later was removed by suction. The oil which was obtained showed no visible change after treatment with 2 cc. of 30% hydrogen peroxide. It was not analyzed further.

¹Farmer and Scott, op. cit.

CHAPTER III

SUMMARY

1. A brief resume of Thiele's theory has been given.
2. It has been shown that Thiele's theory is not universal in application.
3. A brief review of a part of Ingold's theory of addition to unsaturation has been given.
4. The preparation of para-methoxy-diphenylbutadiene has been reported.
5. The preparation of a maleic anhydride derivative of para-methoxy-diphenylbutadiene has also been reported.
6. It has been shown that the bromides of para-methoxy-diphenylbutadiene are unstable in chloroform even in the cold.
7. Straus' prediction concerning the effect of the methoxy-group on unsaturated compounds has been supported.

CHAPTER IV

BIBLIOGRAPHY

Books

- Cohen, J. B. Organic Chemistry for Advanced Students, Vol. I. New York: Longman and Green, and Co., 1928.
- Gilman, H. Organic Chemistry, Vol. I. New York: John Wiley and Sons, Inc., 1938.
- Hilbron, I. Dictionary of Organic Compounds, Vols. I-III. New York: Oxford University Press, 1934.
- Robinson, J. R. Organic Synthesis, Vol. XVI. New York: John Wiley and Sons, Inc., 1935.
- Waters, W. A. Physical Aspects of Organic Chemistry. New York: D. Van Nostrand Co., 1936.
- Watson, H. S. Modern Theories of Organic Chemistry. Oxford: Clarendon Press, 1937.

Periodicals

- Allen, C. F. H. "The Diene Synthesis of Diels and Alder," Journal of Chemical Education, Vol. 10 (1933), pp. 494-497.
- Bergmann, F., and Weinberger, Z. "Remarks on the Synthesis of Substituted Stilbenes and ortho-Phenylbutadienes," Journal of Organic Chemistry, Vol. 6 (1941), pp. 134-39.
- Cain, J. C., Simonsen, J. L., and Smith, C. "Synthesis of Unsymmetrical Derivatives of Deoxybenzoin," Journal of Chemical Society, Vol. 103 (1919), pp. 1035-1039.
- Davies, H., and Davies, W. "The Bromination of Meta-Methoxy Cinnamic Acid," Journal of the Chemical Society, part I (1928), pp. 602-605.
- Farmer, E. H., Lawrence, C. D., and Thorpe, J. F. "Properties of Conjugated Compounds, Part IV. The Formation of Isomeric Additive Dibromides from Butadiene," Journal of the Chemical Society, Part I (1928), pp. 729-739.
- Farmer, E. H., and Scott, W. D. "Properties of Conjugated Compounds, Part VI. The Dibromination Products of Cyclic Butadienes," Journal of the Chemical Society, Part I (1929), pp. 172-180.
- Francis, A. W. "Active Agents in Bromination," Journal of the American Chemical Society, Vol. 47 (1925), pp. 2340-2347.
- Hanson, N. W. and Williams, D. M. "The Addition of Halogen to Unsaturated Acids and Esters. Part III. The Velocity of Bromine Addition," Journal of the Chemical Society,

Part I (1930) pp. 1059-1063.

- Ingold, C. K., and Burton, H. "Mobile Ion Tautomerism, Part I. A Preliminary Study of Conditions of Activation of the Three Carbon System and a Discussion of Results in Relation to Modes of Addition to Conjugated Systems," Journal of The Chemical Society, Part I (1928), pp. 904-921.
- Ingold, C. K., and Smith, H. E. "Mode of Addition to Conjugated Unsaturated Systems. Part III. Further Remarks and Observations Bearing on the Mechanism of Addition of Halogens and Hydrogen Halides," Journal of the Chemical Society, Part II (1931), pp. 2752-2765.
- Meerwein, H., Buchner, E., and Emster, K. van. "Reaction of Aromatic Diazo Compounds upon α , β -Unsaturated Carbonyl Compounds," Journal fur Praktische Chemie, Vol. 152 (1939), pp. 237-266.
- Muskat, I. E., and Huggins, K. A. "Studies of Conjugated Systems. I. The Chlorination of Phenylbutadiene," Journal of the American Chemical Society, Vol. 51 (1929), pp. 2496-2503.
- Riemer, M., and Tobin, E. "The Bromination of 2,4-Dimethoxycinnamic acid," Journal of the American Chemical Society, Vol. 52 (1930), pp. 341-347.
- Riiber, C. N. "Ueber die Constitution des Bisphenylbutadienes," Berichte Der Deutschen Chemischen Gesellschaft, Vol. 42 (1909), pp. 2866-2885.
- Terry, E. M., and Eichelberger, L. "Halogenation of Fumaric and Maleic acids -- Walden's Inversion," Journal of the American Chemical Society, Vol. 47 (1925), pp. 1067-1078.

Unpublished Material

- Proctor, C. A. "The Preparation and Chlorination of 1-p-Chloro-phenyl-4-Phenylbutadiene." Unpublished Master's Thesis, Department of Chemistry, Atlanta University, 1932.
- Yokley, O. E. "Studies in Conjugated Systems: The Addition of Bromine and Hydrogen Bromide to 1-p-Bromo-phenyl-4-Phenyl-Butadiene." Unpublished Master's Thesis, Department of Chemistry, Atlanta University, 1939.